

LA-UR-13-24323

Approved for public release; distribution is unlimited.

Title: Neutron Beta Decay as a Probe of Weak Interactions

Author(s): Saunders, Alexander

Intended for: International Nuclear Physics Conference, 2013-06-01 (Florence, Italy)

Issued: 2013-06-13



Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.

# Neutron Beta Decay as a Probe of Weak Interactions

Alexander Saunders  
Los Alamos National Lab

# Outline

- Why study weak interactions using neutron beta decay?
  - Because of the scientific reach
  - Because of the impact across nuclear and particle physics
- Neutron beta decay correlations
  - How and why you measure them
  - Some sample experiments from the field
- Neutron lifetime
  - The problem and its impact
  - Some recent and upcoming experiments

# Neutron beta decay can have remarkable scientific reach


- Two choices for detecting a force carrier of mass  $M$ :
  - Direct detection, if enough energy available
  - Virtual effects, if not
- If energy is close to mass, then signal may be large and function of energy
- But if energy is much less than mass, then signal is *independent* of energy!
- So the smart thing to do is choose the experiment that gives the best precision

$$S \propto \frac{1}{M^2 - (p/c)^2}$$

$$S \propto \frac{1}{M^2}$$

# Neutron Decay Parameters

- Semi-leptonic decay
  - Lifetime 880 s
  - Endpoint energy 782 keV
- Just two free parameters in SM
  - CKM mixing matrix element
  - Ratio of weak coupling constants
  - Uncertainty comes from radiative corrections



$$n \rightarrow p + e^{-} + \bar{\nu}_e$$



$$\tau_n = \frac{4908.7 \pm 1.9 \text{ s}}{|V_{ud}|^2 (1 + 3\lambda^2)}$$

$$\lambda = g_A / g_V$$


# Neutron beta decay can inform many areas of physics

- Many reactions share the same Feynman diagram as neutron beta decay


Primordial element formation  
( $^2\text{H}$ ,  $^3\text{He}$ ,  $^4\text{He}$ ,  $^7\text{Li}$ , ...)




Solar cycle



Neutron star formation




Pion decay





Neutrino detectors



Neutrino forward scattering




W and Z production



# Decay Correlations


- A: electron asymmetry  $\vec{\sigma}_n \cdot \vec{p}_e$ 
  - Perkeo II, Perkeo III, UCNA
- B: neutrino asymmetry  $\vec{\sigma}_n \cdot \vec{p}_\nu$ 
  - Perkeo II
- C: proton asymmetry  $\vec{\sigma}_n \cdot \vec{p}_p$ 
  - Perkeo II
- D: triple correlation  $\vec{\sigma}_n \cdot (\vec{p}_e \times \vec{p}_\nu)$ 
  - TRINE, emiT
- a: electron-neutrino correlation  $\vec{p}_e \cdot \vec{p}_\nu$ 
  - aSpect, aCORN, Nab



# Search for new physics


- A single parameter yields  $\lambda$ , multiple measurements yield  $V_{ud}$  and beyond
- CKM unitarity  $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1$ 
  - Do neutrons and superallowed beta decays agree?
- Search for right-handed currents (250 GeV limit from n decay)
- Scalar and tensor couplings from  $B$  and  $b$ 
  - Cirigliano 2012

Holeczek *et al.*, arxiv 1303.5295 (2013)






# Principle of the $A$ -coefficient Measurement (and $B$ and $C$ as well)



$$dW = [1 + \beta P A \cos \theta] d\Gamma(E)$$



$$A_{\text{exp}}(E) = \frac{N_1(E) - N_2(E)}{N_1(E) + N_2(E)} \approx \langle P \rangle A \beta \langle \cos \theta \rangle$$

(End point energy = 782 keV)



# Potential Sources of Systematics

- Neutron polarization determination
- Background
  - Due to neutrons that do not decay
- Detector-related effects
  - Electron backscattering and spectroscopy
  - Edge effect (fiducial volume selection)
  - Detector response, calibration
- Statistics
  - Large decay volume, high density

# Perkeo II Experiment Measured ABC


## PERKEO II:

$\beta$ -Asymmetrie	$A = -0.1200 \pm 0.0006$ ,	PRL, in print (2013)
$p$ -Asymmetrie	$B = +0.983 \pm 0.005$ ,	PRL <b>99</b> , 191803 (2007)
$\nu$ -Asymmetrie	$C = -0.239 \pm 0.003$ ,	PRL <b>100</b> , 151801 (2008)




# Perkeo III is state of art of CN beta decay

- Backgrounds eliminated using pulsed beam
- 50 kHz decay rate
- Total uncertainty expected to be  $\Delta A/A = 2.1 \times 10^{-3}$
- Results very soon!




# PERC is the next generation

- **P**roton **E**lectron **R**adiation **C**hannel
- 8 m flight path maximizes statistics
- 6 T field pinch minimizes backscatter, field inhomogeneity effects
- To be installed in flight path at FRM-2
- All systematics expected to be  $O(10^{-4})$



# Interlude — What are ultracold neutrons?

- Very slow neutrons ( $v < 8\text{m/s}$ )
- Totally reflected by some materials
- Hence, they can be **totally confined** within a bottle for periods **in excess of 100 seconds**.
- Typically: **velocity  $< 8\text{m/s}$**   
**kinetic energy  $< 3 \times 10^{-7} \text{ eV}$**   
**wavelength  $> 500\text{\AA}$**   
**or temperature  $< 4 \text{ mK}$**
- *cf:* Gravity:  $10^{-7} \text{ eV/ meter}$ .  
Magnetic field ( $\mu\text{B}$ ):  $10^{-7} \text{ eV/ } 1.7 \text{ T}$ .



# UCNA Experiment — General Approach

Novel features: UCN from pulsed spallation source  
MWPC + plastic scintillator as  $\beta$  detector  
Ultimate Goal: 0.2% measurement of  $A$  ( $\delta A/A = 0.2\%$ )

- Neutron Polarization

- UCN (can produce >99% polarization with 7T magnetic field)
- Copper and diamond-like carbon coated neutron guide (low depolarization)

- Background


- Pulsed UCN source
- MWPC+Plastic scintillator

- Electron backscattering


- MWPC+Plastic scintillator

- Fiducial volume selection

- MWPC





# UCNA Experiment — Apparatus






# UCNA Status

- Taking production data since 2009 using LANL UCN source
- Latest result:  $A = -0.11954(112)$



# Nab experiment in design stage


- Nab will measure  $a, b$  at SNS in US
  - Recall  $a$ =electron-neutrino correlation
  - Reconstruct opening angle from  $E_p, E_e$
  - $E_e$  from Si detectors,  $E_p$  from TOF



# Neutron Lifetime affects BBN

Light elements from  $^2\text{H}$  up to  $^7\text{Li}$  created in "first three minutes"

Weak reactions between particles:



At time  $\approx 1$  s, at "freeze-out" temperature  $T_f \approx 1$  MeV:

neutron to proton ratio frozen to:


$$n/p = \exp(-\Delta m/kT_f) \approx 1/6.$$

$$\Delta m = m_n - m_p = 1.3 \text{ MeV}$$

After another  $\approx 150$  s, practically all neutron wind up in  $^4\text{He}$ ,

i.e., He mass fraction  $Y_p = 2 \times \text{neutron mass fraction} \approx 25\%$ .

Primordial nucleosynthesis:



Dubbers 2013

And ratio of He/n depends directly on n lifetime: 1% lifetime

uncertainty shifts calculated He fraction of  $Y=0.2480 \pm 0.0003$

by 0.0015, or 5 sigma!

# Lifetime uncertainty has grown recently

And central value has shifted by almost 1%!

PDG 2001-2010  $\tau_n = (885.7 \pm 0.8) \text{ s } (S=1)$   
 Serebrov *et al.* 2005  $\tau_n = (878.5 \pm 0.8) \text{ s}$   
 Pichlmaier *et al.* 2010  $\tau_n = (880.7 \pm 1.8) \text{ s}$   
 PDG 2012  $\tau_n = (880.1 \pm 1.1) \text{ s } (S=1.8)$

Beam Method: Count the dying

PHYSICAL REVIEW C **71**, 055502 (2005)





FIG. 2. Experimental method for measuring lifetime by counting neutrons and trapped protons.

**Danger: absolute monitor efficiency needed!**

7 June 2013  
 INPC, Firenze

Serebrov *et al.*,  
 PL B **605**, 72 (2005)  
 Pichlmaier *et al.*,  
 PL B **693**, 221 (2010)



Bottle method: count the survivors



$$1/\tau_{\text{storage}} = 1/\tau_n + 1/\tau_{\text{loss}}$$

Wietfeldt and Greene,  
 Rev. Mod. Phys. **83**, 1173 (2011)

# Material bottle experiments involved 100 s extrapolations due to wall losses



# Solution: eliminate wall losses using magnetic bottle

- A new crop of experiments using magnetic traps is now under development
- Stern-Gerlach effect repels polarized neutrons from walls


ILL Ezhov Bottle filled with vacuum

NIST UCN trap filled with superfluid 4He



increase storage volume from 3.6 l to **15 l**


P. Geltenbort (V. Ezhov)



Top view of the storage bottle made of permanent magnets.

Universidad Autónoma, Madrid, 30 November 2007

45




7 June 2013  
INPC, Firenze

Both have acquired commissioning data;  
HOPE at Munich also in design stage



# Penelope experiment under development


- Superconducting multipole
  - Field zero in center eliminated by inner conductors
- Filled with UCN from FRM-2 through gap in bottom
- Decay products detected at top, guided by field lines
- Spectrum cleaned using absorber lowered from top
- Magnet now under construction




# An outstanding problem: phase space evolution

- Neutron losses on scale of neutron lifetime (quasibound orbits)
- Detector efficiency changes with time
- Must fill phase space evenly, quickly: chaos!

Symmetric Trap has stable orbits      Asymmetric Trap has chaotic orbits



Thin stochastic regions bounded by KAM curves.




Strong perturbation, KAM curves are destroyed.



# UCNTau experiment designed to overcome phase space issues


## Asymmetric Trap $\rightarrow$ Phase Space Mixing

- **Low symmetry** (together with **field ripples**) induces states mixing between circular orbits, through chaotic motion (or not).
- $\rightarrow$  **quick cleaning** ( $\sim$  seconds) of the quasi-bound UCN with large tangential velocities.




# UCNTau Experiment now commissioning

- Asymmetric magnets
- Holding polarization
- UCNs from filled traps
- Neutron absorption inserted activation
- Has stored first time



# New beam-type experiment also nearing first results at J-Parc



- TPC detects electrons from n decay and protons from n-absorption on  ${}^3\text{He}$  simultaneously
- Leading systematic uncertainty expected to be pressure of  ${}^3\text{He}$
- Physics run planned for 2013

# Conclusions

- Precision neutron beta decay measurements can search for physics beyond the standard model with a reach competitive with HEP
- And also underlie wide areas of nuclear and particle physics
- Beta decay correlation experiments in progress and next generation in development: overconstrained SM parameters allow search for BSM effects
- Neutron lifetime problem needs experimental resolution: it is coming, with new experiments worldwide approaching production data
- Review papers:
  - Dubbers and Schmidt, Rev. Mod. Phys. **83**, 1111 (2011)
  - Abele, Pro. Part. Nucl. Phys. **60**, 1 (2008)
  - Wietfeldt and Greene, Rev. Mod. Phys. **83**, 1173 (2011)
- Thank you to my collaborators and especially Dirk Dubbers, Bastian Maerkisch, Hartmut Abele, Chen-Yu Liu, Albert Young